

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/25402>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

## Cardiovascular effects of sulphonylurea derivatives

P. Smits

Departments of Pharmacology and Medicine, University of Nijmegen, Nijmegen, The Netherlands

The classical sulphonylurea derivative glibenclamide is widely prescribed for the treatment of non-insulin-dependent diabetes mellitus (NIDDM). The mechanism of action of this drug is based on an augmented release of insulin from the beta cells of the pancreas, triggered by the closing of so-called ATP-dependent potassium channels (ATP: adenosine'5-triphosphate) [1]. This ultimately results in an influx of calcium via opening of voltage-dependent calcium-ion channels, and the subsequent binding of calcium ions to calmodulin triggers the exocytosis of insulin-containing secretory granules. The central issue of this presentation is the fact that the occurrence of  $K_{ATP}$ -channels is not restricted to the pancreatic beta cells. These particular ion channels occur in several organ systems, including the cardiovascular system [2, 3].

The open state probability of  $K_{ATP}$ -channels is dependent on the cytosolic concentration of ATP. During hypoxia and/or ischaemia, the cytosolic concentration of ATP falls which directly results in opening of  $K_{ATP}$ -channels. The subsequent efflux of potassium induces hyperpolarization of the cell membrane which shortens the action potential in the myocardial cell leading to a reduction in the contractile amplitude of myocardial cells. In vascular smooth muscle cells, opening of  $K_{ATP}$ -channels and subsequent hyperpolarization exerts relaxation of these cells, resulting in an obvious vasodilator response.

The vascular smooth muscle cell  $K_{ATP}$ -channels can be opened pharmacologically by potassium-channel-opening drugs such as cromakalim and the classical vasodilator drug diazoxide [3]. Glibenclamide is able to reduce the vasodilator response to these  $K_{ATP}$  channel-opening drugs, indicating the ability to block vascular  $K_{ATP}$ -channels. Apart from the

interaction with  $K_{ATP}$ -channel-opening drugs, glibenclamide has also been reported to inhibit the vasodilator response to adenosine, and to (endothelium-dependent) vasodilators such as acetylcholine, vasoactive intestinal peptide and insulin [4]. Also in the myocardial cell, glibenclamide is able to reduce the response to  $K_{ATP}$ -channel-opening agents [4].

Animal experiments have shown that the ischaemia-induced opening of myocardial  $K_{ATP}$ -channels is an endogenous cardioprotective mechanism against ischaemia and reperfusion damage. In several ischaemia models glibenclamide increased infarct size [5, 6]. On the other hand blockade of the sulphonylurea receptor is associated with a reduction in ischaemia-related arrhythmias [4]. Not only myocardial  $K_{ATP}$ -channels are involved in the response to ischaemia. Other investigators have demonstrated that the vascular smooth muscle cell response to ischaemia is also inhibited by sulphonylurea derivatives, represented by a reduction of the post-occlusive hyperaemic response after pretreatment with glibenclamide.

Interestingly, the newly developed sulphonylurea derivative glimepiride seems to act more selectively, since its hypoglycaemic action is more pronounced than that of glibenclamide, whereas animal data suggest the absence of interaction with vascular  $K_{ATP}$ -channels [4].

### *Experimental data in humans*

For several reasons the conclusions from animal experiments are not necessarily relevant to the clinical situation for treatment of non-insulin-dependent diabetes (NIDDM) by daily glibenclamide ingestion: 1) the sulphonylurea concentrations used in most animal studies are much higher than those reached during the treatment of NIDDM; 2) in human in vivo

*Corresponding author:* Professor P. Smits, Department of Pharmacology, University of Nijmegen, PO Box 9101, NL-6500 HB Nijmegen, The Netherlands

conditions, glibenclamide is highly protein bound which considerably reduces the free drug concentration; 3) there are large differences between species, and therefore the principles of interaction between glibenclamide and cardiovascular  $K_{ATP}$ -channels are not necessarily relevant to the human situation; and 4) the majority of the experiments show an acute interaction between sulphonylurea derivatives and cardiovascular  $K_{ATP}$ -channels, but this may not be relevant to the chronic setting of NIDDM treatment.

Up to now, human data on this issue are scarce. Kosmas et al. [7] recently showed a reduced post-ischaemic vasodilation after the oral administration of glibenclamide. We recently reported the effects of glibenclamide and tolbutamide on vascular  $K_{ATP}$ -channels in humans [8–10]. In that study, the perfused forearm technique was used. The interaction with vascular  $K_{ATP}$ -channels was studied by investigating the interaction with diazoxide. Therefore, diazoxide was used as a pharmacological tool to open vascular  $K_{ATP}$ -channels. In these studies we showed that glibenclamide was able to attenuate diazoxide-induced and also ischaemia-induced vasodilation. Since the systemic glibenclamide concentrations remained low, our data were not confounded by metabolic effects such as hyperinsulinaemia or hypoglycaemia. Apparently, the perfused forearm technique with infusions of diazoxide as a  $K_{ATP}$ -channel-opener appeared to be an appropriate human in vivo model to quantify the  $K_{ATP}$ -channel-blocking properties of sulphonylurea derivatives at the level of the vascular wall. Therefore, we also tested the newly developed sulphonylurea derivative glimepiride, that has been reported to be devoid of interactions with vascular  $K_{ATP}$ -channels in an animal model. Indeed, when we infused glimepiride into the brachial artery, diazoxide responses were not inhibited. Apparently, pancreatic and vascular  $K_{ATP}$ -channels have different characteristics in humans, which agrees with recent molecular biological observations [8]. Finally, experiments in patients with coronary artery disease have shown that ischaemic preconditioning is inhibited 1 h after the ingestion of 10 mg of glibenclamide [11]. This, as well as the antiarrhythmic properties of glibenclamide in ischaemic conditions, argues for an

interaction between glibenclamide and myocardial  $K_{ATP}$ -channels.

Although the aforementioned data point towards a significant interaction between sulphonylurea derivatives and vascular  $K_{ATP}$ -channels, the clinical relevance of these observations remains unclear. It is obvious that results from clinical trials with well defined end points, like for example the United Kingdom Prospective Diabetes Study (UKPDS), are needed to elucidate this interesting issue.

## References

1. Groop LC (1992) Sulfonylureas in NIDDM. *Diabetes Care* 15: 737–754
2. Noma A (1983) ATP-regulated K-channels in cardiac muscle. *Nature* 305: 147–148
3. Edwards G, Weston AH (1990) Potassium channel openers and vascular smooth muscle relaxation. *Pharmac Ther* 48: 237–258
4. Smits P, Thien Th (1995) Cardiovascular effects of sulphonylurea derivatives; implications for the treatment of NIDDM? *Diabetologia* 38: 116–121
5. Cole WC, McPherson CD, Sontag D (1991) ATP-regulated  $K^+$ -channels protect the myocardium against ischemia/reperfusion damage. *Circ Res* 69: 571–581
6. Gross GJ, Auchampach JA (1992) Blockade of ATP-sensitive potassium channels prevents myocardial preconditioning in dogs. *Circ Res* 70: 223–233
7. Kosmas EN, Levy RD, Hussain SNA (1995) Acute effects of glyburide on the regulation of peripheral blood flow in normal humans. *Eur J Pharmacol* 274: 193–199
8. Bijlstra PJ, Lutterman JA, Russel FGM, Thien Th, Smits P (1996) Interaction of sulphonylurea derivatives with vascular ATP-sensitive potassium channels in humans. *Diabetologia* 39: 1083–1090
9. Bijlstra PJ, Russel FGM, Thien Th, Lutterman JA, Smits P (1996) Effects of tolbutamide on vascular ATP-sensitive potassium channels in humans. *Horm Met Res* 28: 512–516
10. Banitt PF, Smits P, Williams SB, Ganz P, Creager MA (1996) Activation of ATP-sensitive potassium channels contributes to reactive hyperemia in humans. *Am J Physiol* 271: H1594–H1598
11. Tomai F, Crea F, Gasparone A et al. (1994) Ischemic preconditioning during coronary angioplasty is prevented by glibenclamide, a selective ATP-sensitive K channel blocker. *Circulation* 90: 700–705